

FC Professional Insect Control™ Ant Killer Bait Gel

FOR USE IN COMMERCIAL. INDUSTRIAL AND RESIDENTIAL AREAS

ACTIVE INGREDIENT:

OTHER INGREDIENTS:99.999%

†CAS NO. 120068-37-3

Contains: 4 plastic reservoirs

KEEP OUT OF REACH OF CHILDREN CAUTION: SEE CAUTIONS BELOW

Use only in areas not easily accessible to children and pets.

Product may stain textiles and clothing.

This product provides FAST CONTROL of ants in 3-5 days. MAXFORCE® FC PROFESSIONAL INSECT CONTROL™ ANT KILLER BAIT GEL is formulated to effectively control ants. That's because this product contains Fipronil, a fast-acting active ingredient. Its speed is still slow enough, however, for foraging ants to return to the colony and feed the bait to larvae and queen(s) so that the entire colony is killed through the Domino

This product is formulated for effective ant control. The high moisture content of this gel enhances the bait's attractiveness to target ants. The high mosture content of this gel enhances the bait's attractiveness is greatest within two weeks of application. This product is a bait product, formulated to take advantage of ant trophallaxis. Through trophallaxis, worker ants eat the bait while foraging, then feed it to the rest of the colony. The larvae digest the solid bait and produce a liquid containing the Fipronil that is fed to the rest of the colony and

MAXFORCE FC Professional Insect Control Ant Killer Bait Gel with Fipronil effective against: Argentine, Odorous House, Ghost, Pavement, Pharaoh and Little Black Ants

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid contact with ski in and clothing. K Wash hands ther exposed gel away from open foods and food contact surfaces after handling. **FIRST AID:** If swallowed, drink two glasses of by touching back of throat with finger. Do not induce with finger. and induce vomiting live anything by mouth to an unconscious person. Get medical attention. In or niah 1-800-426-6228

STORAGE AND DISPOSAL: Do not contaminate storage or dispos STORAGE: Store in a cool, dry area out of reach

DISPOSAL: Wrap used prefilled res

For questions or comments about MAXFORCE, the MAXFORCE website at www.naxforce.com. call toll-free 1-800-426-6228

DIRECTIONS FOR USE.

It is a violation of Federa law to use this ro nt with its la uct in a manner in Application rate is dependent on level of in estation and species to be should be applied as spots.

Package should be in possession of applicator.

- 1. Apply gel in spots in cracks and crevices. For best results, place many spots or dabs of gel where you see ants foraging and entering the structure.
- 2. Do not apply gel to areas which have been recently sprayed with insecticide or contaminated with foggers, and do not spray insecticide over gel as this may cause the bait to become repellent. Do not apply gel to areas that are routinely washed, such as counter tops, as gel will be removed or diluted and spread onto the surface. If you have recently sprayed, remove the dead ants and thoroughly clean the area before applying ge
- 3. Place the bait as follows: Gel can be placed in cracks, crevices, corners and any out-ofthe-way place, such as under and behind kitchen appliances, along baseboards, under sinks, in pantry, around pipes and water heater, window and door frames, and in the
- two weeks. Inspect gel placements and reapply as necessary. Reapply in same area if bait is gone or in a different location if no ant activity is evident.

Applications in Food/Feed Handling Establishments. Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only.

garage, attic, or crawl space. 4. Inspect placements and reapply as needed: The gel will remain attractive to ants for up to

Apply with syringe applicator or bait injector directly into cracks and crevices. Place applicator tip into cracks and crevices and where ants may travel. Apply gel (up to 1 gram) per spot. Apply bait in small amounts between different elements of construction, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, thors and windows, cabinets, sinks, stoves and other places where ants may enter the establishment. During follow-up visits inspect bait placements and reapply when necessary.

Do not apply bait to areas where food feed, food utensils or food processing surfaces may become contaminated. Do not apply bait in cracks and crevices of areas that are routinely washed food of tabes, to od/feed preparation and prepared food/feed holding surfaces as bait pray be removed by washing. Do not apply bait to surfaces where the temperature exceeds 1207 as bait may rub. Examples include, but are not limited to, portions of stoves, overs, gribs, fur he hoots, heat lamps, coffee urns, steam tables, toasters, fiyers, dishwashers and hot water pipes. Care should be taken to avoid depositing gel onto exposed surfaces. If yell contacts an exposed surface, remove it and wash exposed surface.

eas include areas for receiving, storage, packing (canning, bottling, wrapping,

nod/ned areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), peparing edible waste storage, and enclosed processing systems (mills, dairies, dible bils, syrups). Serving areas are also considered a food area when food is exposed and facility is in operation. Application of this product in food/feed areas of food/feed handling stablishments other than as a crack and crevice treatment is not permitted. Gel placements should be year and trails, next to ant nests or in areas of suspected and activity.

For indoor user Visually inspect for arts and place gel in areas where you see ants. Likely locations for ants and gel placements are areas where ants enter the building structure and areas with potential water damage including: wall voids, cracks and crevices in kitchen, bathroom, under sight, window sills, screen vents, garage, attic, and crawl spaces. Do not place bait to be attions that are routinely washed, as bait will be removed by vastain. Do not treat food preparation surfaces. Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

Vor cutdoor use: Place gel outdoors into, on or adjacent to structures where you see ans, nex to ant trails and suspected areas of ant activity. Likely locations for ants and cements are points of entry into structure and areas with potential water damage including: eaves, soffits, weep holes, where wires and pipes come into the building structure, trees, stumps, roots, wood decks, fences, planter boxes, woodpiles, landscaping, and railroad ties. Extreme heat or cold, direct sunlight and water may educe the residual effectiveness. Apply Gel outdoors into MAXFORCE Refillable Bait Stations or other devices designed to hold and protect bait outdoors. Close Refillable MAXFORCE Bait Station securely after application.

Pest population reduction will be apparent within a week.

Directions for using the Maxforce Bait Injector, loading reservoirs and operation: With the teeth of the drive rod facing up, pull the drive rod all the way back. Attach ant gel reservoir to the bait injector with a twist-locking movement. Remove the reservoir outlet cap and attach the dispensing tip. Push the drive rod until the drive plunger seats against the piston in the reservoir of gel. Turn drive rod knob until teeth are facing down. Gently squeeze the trigger until the gel flows out. Reservoirs that still contain gel can be sealed with the end cap for future use.

Directions for changing reservoirs: With the arrow on the drive rod pointing up, pull the drive rod out as far as it will go. Remove the empty reservoir, wrap, and place in trash can. Reservoirs that still contain gel can be sealed with the outlet cap for future use. To attach a new reservoir of gel, follow the above directions.

> Contains 4 plastic reservoirs NET WEIGHT 0.95 oz (27g) TOTAL NET WEIGHT 3.80 oz. (108 g)

FC Professional Insect Control™Ant Killer Bait Gel

Distributed by & © 2001 Maxforce Insect Control Systems 1221 Broadway Oakland, CA 94612

EPA Reg. No. 64248-21 EPA Est. No. 5813-MS-001